

參考文獻

- [1] Cheng, S. W. and Ye, K. Q. (2004). Geometric Isomorphism and Minimum Aberration for Factorial Designs with Quantitative Factors. *The Annals of Statistics* **32**, 2168-2185.
- [2] Cox, D., Little, J. and O'Shea, D. (2007). *Ideals, Varieties and Algorithms, An Introduction to Computational Algebraic Geometry and Commutative Algebra*, 3rd ed, Springer-Verlag New York, Inc.
- [3] Fontana, R., Pistone, G., Rogantin, M.P. (1997). Algebraic analysis and generation of two-levels designs. *Statist. Appl.* **9**, 15-29.
- [4] Fontana, R., Pistone, G. and Rogantin, M. P. (2000). Classification of two-level factorial fractions. *J. Statist. Plann. Inference* **87** 149–172.
- [5] Pistone, G., Riccomagno, E. and Wynn, H. P. (2001). *Algebraic Statistics: Computational Commutative Algebra in Statistics*. Chapman and Hall, Boca Raton, FL.
- [6] Pistone, G. and Wynn, H.P. (1996). Generalised confounding with Grobner bases. *Biometrika* **83**, 653-666.
- [7] Ye, K. Q. (2003). Indicator Function and its Application in Two-level Factorial Designs. *The Annals of Statistics* **31**, 984-994.