

I. Notations and definitions

In the following problem set, the symbols  $\mathbb{N}$  and  $\mathbb{R}$  will be reserved for the set of all positive integers and the set of all real numbers, respectively.

We shall fix a field  $K$  and denote by  $V, W$  and  $W'$  finite dimensional vector spaces over  $K$  and we shall use  $\dim V$  for the dimension of  $V$ . For any linear transformation  $f : V \rightarrow W$ ,  $\ker f$  is the kernel of  $f$  and  $\text{Im } f$  is the image of  $f$ .

The letters  $n$  and  $r$  will denote natural numbers, and  $\text{Mat}_n(K)$  ( $\text{Mat}_r(K)$ ) will denote the set of all  $n$  by  $n$  ( $r$  by  $r$ , resp.) matrices over  $K$ . We call two matrices  $A, B \in \text{Mat}_n(K)$  similar if there exists an invertible matrix  $P \in \text{Mat}_n(K)$  such that  $P^{-1}AP = B$ . A matrix  $A \in \text{Mat}_n(K)$  is called nilpotent if  $A^s = 0$  for some  $s \in \mathbb{N}$ .

II. Problems

- (1) Let  $E$  be a subset of  $V$  and  $\langle E \rangle = \{ \sum_{\alpha \in K, v \in E} \alpha v \mid \text{only finitely many } \alpha \text{ are nonzero} \}$ . Show that  $\langle E \rangle$  is a subspace of  $V$  and that  $\langle E \rangle = \bigcap \{ W \mid W \text{ is a subspace and } E \subseteq W \}$ . 10%
- (2) Let  $f : V \rightarrow W$  be a linear transformation. Then there exist bases  $\{u_1, \dots, u_m\}$  of  $V$  and  $\{v_1, \dots, v_n\}$  of  $W$  and a positive integer  $r, r \leq m$  and  $r \leq n$ , such that  $f(u_i) = v_i$  for  $i = 1, \dots, r$  and  $f(u_i) = 0$  for  $i = r + 1, \dots, m$ . 10%
- (3) Let  $f : V \rightarrow W$  and  $g : V \rightarrow W'$  be linear transformations such that  $\ker g \subseteq \ker f$ . Show that there exists a linear function  $h : W \rightarrow W'$  such that  $h \circ g = f$ . (Hint. Consider extending a basis of  $\ker g$  to a basis of  $V$  and remember that  $\dim V = \dim(\ker g) + \dim(\text{Im } g)$ .) 15%
- (4) (i) Let  $\{x_1, \dots, x_m\}$  be a basis of  $V$ . Suppose that  $\alpha_1, \dots, \alpha_m \in K$  are pairwise distinct scalars. If  $f : V \rightarrow V$  is a linear transformation such that  $f(x_i) = \alpha_i x_i$  for  $i = 1, 2, \dots, m$ , and if  $g : V \rightarrow V$  is a linear transformation with the property that  $f \circ g = g \circ f$ , then there exist scalars  $\beta_1, \dots, \beta_m \in K$  such that  $g(x_i) = \beta_i x_i$  for  $i = 1, 2, \dots, m$ . 10%  
 (ii) Show that if  $A \in \text{Mat}_m(K)$  and  $AB = BA$  for all  $B \in \text{Mat}_m(K)$ , then  $A$  is a scalar diagonal matrix. 10%  
 (iii) Show that if  $A, B, C \in \text{Mat}_2(K)$ , then  $(AB - BA)^2$  commutes with  $C$ . 5%
- (5) If  $A, B \in \text{Mat}_n(K)$  with  $A$  invertible, then the matrix  $A + rB$  is invertible for all but finite number of  $r \in K$ . 15%
- (6) Let  $A \in \text{Mat}_n(K)$  be nilpotent. Then  $A$  is similar to a matrix of the form 15%

$$\begin{pmatrix} W & 0 \\ 0 & 0 \end{pmatrix},$$

where  $W \in \text{Mat}_r(K), 1 \leq r \leq n$ , is of the form

$$W = \begin{pmatrix} 0 & 1 & 0 & 0 & \dots & 0 \\ 0 & 0 & 1 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \vdots & & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & \dots & 1 \\ 0 & 0 & 0 & 0 & \dots & 0 \end{pmatrix}.$$

- (7) For each of the following matrices  $A$  and  $B$ , determine whether or not it is diagonalizable over  $\mathbb{R}$ . If it is diagonalizable, then find the change-of-coordinate matrix which make it a diagonal matrix, otherwise, explain why it cannot be diagonalized. 10%

$$A = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 2 & 1 \\ 0 & 0 & 3 \end{pmatrix}, \quad B = \begin{pmatrix} 3 & 1 & 0 & 0 \\ 0 & 3 & 0 & 0 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 2 \end{pmatrix}.$$