




# Chap.2 Supplement

*For the EM Course Lectured by Prof. Tsun-Hsu Chang  
Teaching Assistants: Hung-Chun Hsu, Yi-Wen Lin, and Tien-Fu Yang  
2022 Fall at National Tsing Hua University*






# Brief History of Field theory





Issac Newton  
(1643-1727)

- Classical field theory can be traced back to Newton's law of universal gravitation.
- “Action at a distance”:  
**Instantaneous effect** on faraway objects
- Newton couldn't find empirical evidence to support a causal explanation of gravity, and any description remained purely hypothetical.
- Concept of fields was proposed by mathematical physicists in the 18<sup>th</sup> century as a **mathematical trick without physical interpretation**.






# Brief History of Field theory



Michael Faraday  
(1791-1867)

- Michael Faraday coined the term “field” in 1845.
- He stated
  - Interactions between objects occur via **space-filling “lines of force”** that can move, expand, and contract.
  - Atoms are convergences of the lines of force.
  - Non-instantaneous EM interactions are fatal to AD.
  - Magnetic lines of force are physical conditions of **mere space** (space without substance).





# Brief History of Field theory



James Maxwell  
(1831-1879)

- Maxwell was more “Newtonian” and more mathematical than Faraday.
- Faraday’s lines of force is the state of **the ether** as a velocity field is a state of a material fluid.
- His analysis led to the predictions of
  - a **finite transmission time** for EM actions
  - **light** as an EM field phenomenon
- A time delay demands **the presence of energy in the intervening space** to conserve energy.
- Action-at-a-distance was thus conclusively refuted.





# Brief History of Field theory




“Before Maxwell, Physical Reality was thought of as consisting in material particles ... **Since Maxwell’s time, Physical Reality has been thought of as represented by continuous fields**, ... and not capable of any mechanical interpretation. This change in the conception of Reality is the most profound and fruitful that physics has experienced since the time of Newton.”

- Albert Einstein -






# Brief History of Field theory




Discrete lines in atomic spectra



Distribution of blackbody radiation  
in different wavelengths

→ The beginning of quantum mechanics!





# Brief History of Field theory

Max Planck (1858-1947)

Atoms **discretely absorb and emit EM radiation** as tiny oscillators.

→ Quantization



Louis de Broglie (1892-1987)

Microscopic particles exhibit both **wave- and particle-like properties under different circumstances.**

→ Wave-Particle Duality



Werner Heisenberg (1901-1976)

**Matrix formulation** and Law of multiplication (Commutator)

→ Creation of Quantum Mechanics (with help of Born and Jordan)




Albert Einstein (1879-1955)

Light is composed of **individual packets of energy** called photons.

→ Photoelectric effect

The distinction between time and space was blurred.

→ **Special Relativity**





## Two Difficulties Remained

- Schrödinger equation explains stimulated emission but not **spontaneous emission**.
- Schrödinger equation treats **time as an ordinary number while promoting spatial coordinates to linear operators.**



# Brief History of Field theory



Albert Einstein  
(1879-1955)

- Einstein supported a “**fields are all there is**” view of classical physics.

→ **Fields are states of space.**

- The general theory of relativity (GR) resolves Newton’s dilemma about the “absurdity of gravitational Action at a distance”.

$$\underbrace{R_{\mu\nu}(x) - \frac{1}{2}g_{\mu\nu}(x)R(x)}_{\text{Geometry of space-time}} + \underbrace{g_{\mu\nu}(x)\Lambda}_{\text{Energy and momentum of matter}} = kT_{\mu\nu}(x)$$



Geometry of space-time

Energy and momentum  
of matter

**Matter tells space-time how to curve; curved space-time tells matter how to move.**



# Brief History of Field theory







# Brief History of Field theory

Further development of fields leads to the conclusion:

Charges are **emitting particles** constantly, and forces/ interactions result from exchanging particles between charges.




Yukawa (1949 Nobel)





# Brief History of Field theory

## Standard Model of Elementary Particles



The view of gauge bosons is now the basis for “Standard Model” of elementary particles.

- Strong: Gluons
- EM : Photons
- Weak : W, Z bosons
- Gravitational : Graviton? (not observed yet)


Higgs bosons: the mechanism “to get mass”

延伸閱讀：上帝的粒子－希格斯粒子的發明與發現



# Brief History of Field theory

From the above brief introduction, it is clear that **electromagnetism** is the first example of the modern **theory of interaction**. That's why we need **tensor/vector analysis** to describe **fields**.






# Three-body Problem

$$\begin{aligned}\ddot{\mathbf{r}}_1 &= -Gm_2 \frac{\mathbf{r}_1 - \mathbf{r}_2}{|\mathbf{r}_1 - \mathbf{r}_2|^3} - Gm_3 \frac{\mathbf{r}_1 - \mathbf{r}_3}{|\mathbf{r}_1 - \mathbf{r}_3|^3} \\ \ddot{\mathbf{r}}_2 &= -Gm_3 \frac{\mathbf{r}_2 - \mathbf{r}_3}{|\mathbf{r}_2 - \mathbf{r}_3|^3} - Gm_1 \frac{\mathbf{r}_2 - \mathbf{r}_1}{|\mathbf{r}_2 - \mathbf{r}_1|^3} \\ \ddot{\mathbf{r}}_3 &= -Gm_1 \frac{\mathbf{r}_3 - \mathbf{r}_1}{|\mathbf{r}_3 - \mathbf{r}_1|^3} - Gm_2 \frac{\mathbf{r}_3 - \mathbf{r}_2}{|\mathbf{r}_3 - \mathbf{r}_2|^3}\end{aligned}$$


$$\mathcal{H} = -\frac{Gm_1m_2}{|\mathbf{r}_1 - \mathbf{r}_2|} - \frac{Gm_2m_3}{|\mathbf{r}_3 - \mathbf{r}_2|} - \frac{Gm_3m_1}{|\mathbf{r}_3 - \mathbf{r}_1|} + \frac{\mathbf{p}_1^2}{2m_1} + \frac{\mathbf{p}_2^2}{2m_2} + \frac{\mathbf{p}_3^2}{2m_3}$$



Chaos!


So... Can we find an analytic solution by “Field”?





Illustrations: Niklas Elmehed

## THE NOBEL PRIZE IN PHYSICS 2022



Alain  
Aspect

John F.  
Clauser

Anton  
Zeilinger

"for experiments with entangled photons,  
establishing the violation of Bell inequalities  
and pioneering quantum information science"

THE ROYAL SWEDISH ACADEMY OF SCIENCES



# Homework Exercises

Griffiths:

2.5, 2.6, \*2.9, 2.10, 2.14, 2.16, 2.18, \*2.20, 2.21, 2.22, 2.25,  
2.27, 2.29, 2.30, 2.33, 2.34, \*2.36, 2.38, \*2.39, 2.42, \*2.43,  
\*2.53, 2.60

Open question: Is electric/magnetic force generally a conservative force? Why or why not?






# Griffiths Prob 2.53

**Problem 2.53** In a vacuum diode, electrons are “boiled” off a hot **cathode**, at potential zero, and accelerated across a gap to the **anode**, which is held at positive potential  $V_0$ . The cloud of moving electrons within the gap (called **space charge**) quickly builds up to the point where it reduces the field at the surface of the cathode to zero. From then on, a steady current  $I$  flows between the plates.

Suppose the plates are large relative to the separation ( $A \gg d^2$  in Fig. 2.55), so that edge effects can be neglected. Then  $V$ ,  $\rho$ , and  $v$  (the speed of the electrons) are all functions of  $x$  alone.







**FIGURE 2.55**





# Inventor of Light Bulb




Further reading: 怪奇科學研究所: 42個腦洞大開的趣味科學故事




# “Edison” effect

Vacuum diode with forward voltage



Vacuum diode with reverse voltage



Richardson-Dushman Equation

$$J = aT^2 \exp(-W / RT)$$



Edison saw no special value in this effect, but he patented it anyway.  
Today we call the effect by the more descriptive term, “thermionic emission.”



# Further Applications



John Ambrose Fleming  
(1849 ~ 1945)




Single-molecule optoelectronic devices promise a potential solution for miniaturization and functionalization of silicon-based microelectronic circuits in the future.

Ref.: P. Li et al., *Rep. Prog. Phys.* **85**, 086401 (2022).



# Griffiths Prob 2.53

Before the calculation: What is **the physics**? → Dimensional Analysis



Current Density  $\mathbf{J} = \rho \mathbf{v}$  ← Two scales needed

Length:  $d$  (m),  $A$  (m<sup>2</sup>)

Mass:  $m$  (kg)

Charge:  $e$  (A·s)

Voltage:  $V_0$  (kg·m<sup>2</sup>·s<sup>-3</sup>·A<sup>-1</sup>)

Permittivity:  $\epsilon_0$  (kg<sup>-1</sup>·m<sup>-3</sup>·s<sup>4</sup>·A<sup>2</sup>)

For moving electrons in solids: introduce mobility  $\mu$  (response to E field)

Velocity scale  $v_s = \sqrt{eV_0/m}$

Charge Density scale  $\rho_s = \epsilon_0 V_0 / d^2$

$$\therefore J_s \sim v_s \rho_s \sim \epsilon_0 \sqrt{\frac{e}{m}} \frac{V_0^{3/2}}{d^2}$$

Known as Child-Langmuir Law


$$\text{Velocity scale } v_s = \mu \cdot V_0 / d \quad \therefore J_s \sim v_s \rho_s \sim \epsilon_0 \mu^0 \frac{V^2}{d^3} \quad \text{Known as Mott-Gurney law}$$



# Griffiths Prob 2.53

Velocity scale  $v_s = \sqrt{eV_0/m}$

Charge Density scale  $\rho_s = \epsilon_0 V_0 / d^2$



Conservation of energy

$$\frac{1}{2}mv^2(z) - e\phi(z) = \text{Const.} = 0$$

$$J_s \sim v_s \rho_s \sim \epsilon_0 \sqrt{\frac{e}{m}} \frac{V_0^{3/2}}{d^2} \propto \frac{[\phi(z)]^{3/2}}{d^2} \leftarrow \text{Constant for all } z$$

$$\Rightarrow \phi(z) = V_0 \left( \frac{z}{d} \right)^{4/3} \Rightarrow \rho(z) = -\epsilon_0 \cdot \phi'' = \frac{4}{9} \epsilon_0 \frac{V_0}{d^2} \left( \frac{z}{d} \right)^{-2/3}$$


$$\therefore J = -\rho(z)v(z) = \frac{4}{9} \epsilon_0 \frac{V_0}{d^2} \left( \frac{z}{d} \right)^{-2/3} \sqrt{\frac{2e\phi(z)}{m}} = \frac{4}{9} \epsilon_0 \sqrt{\frac{2e}{m}} \frac{V_0^{3/2}}{d^2}$$

$$\Rightarrow I = JA = \frac{4}{9} \epsilon_0 \sqrt{\frac{2e}{m}} V_0^{3/2}$$

Ref.: R. J. Umstadtd et al., *Am. J. Phys.* **73**, 160 (2005).



# Griffiths Prob 2.53



Or you can solve this by brutal force.....

$$\nabla^2 \phi(z) = -\frac{\rho(z)}{\epsilon_0} \quad \frac{1}{2} m v^2(z) - e\phi(z) = 0 \Rightarrow v = \sqrt{\frac{2e\phi}{m}}$$

$$J(z) = \rho v \Rightarrow \rho = J \sqrt{\frac{m}{2e\phi}}$$


B.C.:  $\phi(0) = \phi'(0) = 0, \phi(d) = V_0$

$$\phi' \phi = -\frac{J}{\epsilon_0} \sqrt{\frac{m}{2e}} \frac{1}{\sqrt{\phi}} \Rightarrow \int \phi' d\phi = -\frac{J}{\epsilon_0} \sqrt{\frac{m}{2e}} \int \phi'^{-1/2} dz d\phi'$$

$$\frac{1}{2} \phi'^2 = -\frac{J}{\epsilon_0} \sqrt{\frac{m}{2e}} \phi'^{1/2} \Rightarrow \frac{d\phi}{dz} = \dots \Rightarrow \phi = V_0 \left( \frac{z}{d} \right)^{4/3} \Rightarrow \rho(z) = -\epsilon_0 \cdot \phi' = -\frac{4}{9} \epsilon_0 \frac{V_0}{d^2} \left( \frac{z}{d} \right)^{-2/3}$$



# Griffiths Prob 2.53



$$J = \frac{4}{9} \epsilon_0 \sqrt{\frac{2e}{m}} \frac{V_0^{3/2}}{d^2}$$

Assumptions:

1. Steady-state in the vacuum gap
2. Single species charged particles with zero initial velocity
3. Nonrelativistic and non-quantum under electrostatic approximation.




FIG. 2 (color online). The values of  $\mu_Q V_g^{3/2}$  as a function of gap voltage  $V_g$  (volt) for various gap spacing  $D = 1, 10, \text{ and } 100 \text{ nm}$  (solid lines: top to bottom). The dashed lines are without an exchange-correlation term ( $\phi_{xc} = 0$ ) for  $D = 10$  and  $100 \text{ nm}$  (top to bottom), and the short-dashed line is the classical limit.

Ref.: L. K. Ang, T. J. T. Kwan, and Y. Y. Lau, *Phys. Rev. Lett.*, **91**, (2003).



# Further reading

APPLIED PHYSICS REVIEWS 4, 011304 (2017)



## APPLIED PHYSICS REVIEWS

### 100 years of the physics of diodes

Peng Zhang,<sup>1,a)</sup> Ágúst Valfells,<sup>2</sup> L. K. Ang,<sup>3</sup> J. W. Luginsland,<sup>4</sup> and Y. Y. Lau<sup>1</sup>

<sup>1</sup>Department of Nuclear Engineering and Radiological Sciences, University of Michigan, Ann Arbor, Michigan 48109-2104, USA

<sup>2</sup>School of Science and Engineering, Reykjavik University, Menntavegi 1, 101 Reykjavik, Iceland

<sup>3</sup>Engineering Product Development, Singapore University of Technology and Design, Singapore, Singapore 487372

<sup>4</sup>Air Force Office of Scientific Research, Arlington, Virginia 22203, USA

(Received 29 September 2016; accepted 29 November 2016; published online 17 March 2017)

The Child–Langmuir Law (CL), discovered a century ago, gives the maximum current that can be transported across a planar diode in the steady state. As a quintessential example of the impact of space charge shielding near a charged surface, it is central to the studies of high current diodes, such as high power microwave sources, vacuum microelectronics, electron and ion sources, and high current drivers used in high energy density physics experiments. CL remains a touchstone of fundamental sheath physics, including contemporary studies of nanoscale quantum diodes and nano gap based plasmonic devices. Its solid state analog is the Mott–Gurney law, governing the maximum charge injection in solids, such as organic materials and other dielectrics, which is important to energy devices, such as solar cells and light emitting diodes. This paper reviews the important advances in the physics of diodes since the discovery of CL, including virtual cathode formation and extension of CL to multiple dimensions, to the quantum regime, and to ultrafast processes. We review the influence of magnetic fields, multiple species in bipolar flow, electromagnetic and time dependent effects in both short pulse and high frequency THz limits, and single electron regimes. Transitions from various emission mechanisms (thermionic-, field-, and photoemission) to the space charge limited state (CL) will be addressed, especially highlighting the important simulation and experimental developments in selected contemporary areas of study. We stress the fundamental physical links between the physics of beams to limiting currents in other areas, such as low temperature plasmas, laser plasmas, and space propulsion.

[<http://dx.doi.org/10.1063/1.4978231>]

P. Zhang, Á. Valfells, L. K. Ang, J. W. Luginsland, and Y. Y. Lau, *Appl. Phys. Rev.* 4, 011304 (2017).

