




Quiz 3 (Chap. 4)

*For the EM Course Lectured by Prof. Tsun-Hsu Chang
Teaching Assistants: Hung-Chun Hsu, Yi-Wen Lin, and Tien-Fu Yang
2022 Fall at National Tsing Hua University*







What did we learn in Chap. 4?

Conductors vs. dielectrics (or insulators)

Polar



Non-Polar




The atom or molecule now has a tiny dipole moment $\mathbf{p}$, which **points in the same direction as $\mathbf{E}$** and is proportional to the field.

$$\mathbf{p} = \alpha_{ij} \mathbf{E}, \quad \alpha = \text{atomic polarizability}$$

Rule of Thumb! Not a fundamental Law!

α_{ij} : polarizability tensor for the molecule

Always possible to choose “principal” axes such that the off-diagonal terms vanish, leaving just three nonzero polarizabilities.





What did we learn in Chap. 4?

Example 4.1 A primitive model for an atom consists of a point nucleus (+q) surrounded by a uniformly charged spherical cloud (−q) of radius a.




Figure 4.1




Figure 4.2


$$E_e = \frac{1}{4\pi\epsilon_0} \frac{qd}{a^3} \quad p = qd = (4\pi\epsilon_0 a^3)E = \alpha E$$

$\alpha = 4\pi\epsilon_0 a^3$ the atomic polarizability

Prob.4.2 According to **quantum mechanics**, the electron cloud for a hydrogen atom in ground state has a charge density

$$\rho(r) = \frac{q}{\pi a^3} e^{-2r/a},$$

$$E_{e,QM}(\mathbf{r}) \sim \frac{qd}{3\pi\epsilon_0 a^3} \sim \frac{p}{\alpha}$$



$$\left\{ \begin{array}{l} \frac{\alpha_{qm}}{4\pi\epsilon_0} \sim \frac{3}{4} a^3 \sim 0.09 \times 10^{-30} m^3 \\ \frac{\alpha_{cl}}{4\pi\epsilon_0} \sim a^3 \sim 0.12 \times 10^{-30} m^3 \end{array} \right.$$

Experiment
Hydrogen atom $\sim 0.667 \times 10^{-30}$



What did we learn in Chap. 4?


Example 4.3 If we have two spheres of charge: a positive sphere and a negative sphere. When the material is uniformly polarized, all the plus charges move slightly upward (the z-direction), all the minus charges move slightly downward. The two spheres no longer overlap perfectly. Find the polarizability.

Sol. The electric field inside a uniform charged sphere of radius a


$$\mathbf{E}_e(r) = \frac{1}{4\pi r^2} \frac{\frac{4}{3}\pi r^3 \rho}{\epsilon_0} \hat{\mathbf{r}} = \frac{\rho r}{3\epsilon_0} \hat{\mathbf{r}} = \frac{1}{4\pi\epsilon_0} \frac{qr}{a^3} \hat{\mathbf{r}}, \text{ where } q = \frac{4}{3}\pi a^3 \rho$$

Two uniformly charged spheres separated by $\mathbf{d}$ produce the electric field:

$$\mathbf{E}(\mathbf{r}) = \mathbf{E}_{q+}(\mathbf{r}_+) + \mathbf{E}_{q-}(\mathbf{r}_-) = \frac{1}{4\pi\epsilon_0} \frac{q}{a^3} (\mathbf{r}_+ - \mathbf{r}_-) = \frac{1}{4\pi\epsilon_0} \frac{q}{a^3} \left(\left(\mathbf{r} - \frac{1}{2}\mathbf{d} \right) - \left(\mathbf{r} + \frac{1}{2}\mathbf{d} \right) \right) = -\frac{1}{4\pi\epsilon_0} \frac{q\mathbf{d}}{a^3} = -\frac{1}{4\pi\epsilon_0} \frac{q^2}{a^3} \mathbf{p}$$



$$V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \frac{\hat{\mathbf{r}} \cdot \mathbf{p}}{r^2}$$






What did we learn in Chap. 4?

Example 3.10 An electric dipole consists of two equal and opposite charges separated by a distance d . Find the approximate potential V at points far from the dipole.

$$V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \frac{qd \cos\theta}{r^2} = \frac{p}{4\pi\epsilon_0} \frac{\cos\theta}{r^2} = \frac{1}{4\pi\epsilon_0} \frac{\hat{\mathbf{r}} \cdot \mathbf{p}}{r^2}$$

where $\mathbf{p} = q\mathbf{d}$ pointing from negative charge to positive charge.






What did we learn in Chap. 4?

In a uniform field, the force on the positive end, $\mathbf{F} = q\mathbf{E}$, exactly cancels the force on the negative end. However, there will be a torque:

$$\mathbf{N} = (\mathbf{r}_+ \times \mathbf{F}_+) + (\mathbf{r}_- \times \mathbf{F}_-) = q\mathbf{d} \times \mathbf{E} = \mathbf{p} \times \mathbf{E}$$

in such a direction as to line $\mathbf{p}$ up parallel to $\mathbf{E}$



The force on a dipole in a **non**uniform field

$$\mathbf{F} = q(\mathbf{E}_+ - \mathbf{E}_-) \cong q((\mathbf{d} \cdot \nabla)\mathbf{E}) \cong (\mathbf{p} \cdot \nabla)\mathbf{E}$$

What happens to a piece of dielectric material when it is placed in an electric field?

A lot of little dipoles point along the direction of the field and the material becomes polarized.

A convenient **measure of this effect is $\mathbf{P} \equiv$ dipole moment per unit volume**, which is called the Polarization.



What did we learn in Chap. 4?

$$V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_v \frac{\hat{\mathbf{r}} \cdot d\mathbf{p}}{r^2} = \frac{1}{4\pi\epsilon_0} \int_v \frac{\hat{\mathbf{r}} \cdot \mathbf{P}(\mathbf{r}')}{r^2} d\tau' = \frac{1}{4\pi\epsilon_0} \int_v \mathbf{P} \cdot \nabla' \left(\frac{1}{r} \right) d\tau'$$


$$V = \frac{1}{4\pi\epsilon_0} \left[\int_v \nabla' \cdot \left(\frac{\mathbf{P}}{r} \right) d\tau' - \int_v \frac{1}{r} (\nabla' \cdot \mathbf{P}) d\tau' \right]$$

$$= \frac{1}{4\pi\epsilon_0} \oint_s \frac{\mathbf{P}}{r} \cdot d\mathbf{a}' + \frac{1}{4\pi\epsilon_0} \int_v \frac{1}{r} (-\nabla' \cdot \mathbf{P}) d\tau'$$

$$\begin{cases} \sigma_b = \mathbf{P} \cdot \hat{\mathbf{n}} \\ \rho_b = -\nabla' \cdot \mathbf{P} \end{cases}$$

$$V = \frac{1}{4\pi\epsilon_0} \oint_s \frac{\sigma_b}{r} da' + \frac{1}{4\pi\epsilon_0} \int_v \frac{\rho_b}{r} d\tau'$$


**The potential of a polarized object
= produced by a surface charge density
+ a volume charge density**





What did we learn in Chap. 4?

Ex. 4.2 Find the electric field produced by a uniformly polarized sphere of radius R .




$$\begin{cases} \sigma_b = \mathbf{P} \cdot \hat{\mathbf{n}} = P \cos \theta' \\ \rho_b = -\nabla' \cdot \mathbf{P} = 0 \end{cases}$$

$$V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_0^\pi \frac{P \cos \theta'}{r} 2\pi R^2 \sin \theta' d\theta'$$

$$V(r, \theta, 0) = \begin{cases} \frac{1}{3\epsilon_0} \frac{PR^3}{r^2} \cos \theta & (r \geq R) \\ \frac{P}{3\epsilon_0} r \cos \theta & (r \leq R) \end{cases}$$

$$\begin{aligned} \frac{1}{r} &= \frac{1}{r} \sum_{n=0}^{\infty} \left(\frac{R}{r}\right)^n P_n(\cos \theta') & r \geq R \\ &= \frac{1}{R} \sum_{n=0}^{\infty} \left(\frac{r}{R}\right)^n P_n(\cos \theta') & r \leq R \end{aligned}$$

$\therefore$ orthogonality $\therefore$ only $n=1$ survive



$$\mathbf{E} = -\nabla V = -\frac{P}{3\epsilon_0} \hat{\mathbf{z}} \quad \text{uniformly}$$





What did we learn in Chap. 4?

The electric field inside matter

Microscopic level → Too Complicated to calculate

Macroscopic field → Defined as the average field over regions large enough

For many substances, the polarization is proportional to the field, provided $\mathbf{E}$ is not too strong.

$$\mathbf{P} = \epsilon_0 \chi_e \mathbf{E} \quad \chi_e : \text{the electric susceptibility}$$

The total field $\mathbf{E}$ may be due in part to **free charges** and in part to **the polarization itself**.


Materials that obey above equation are called linear dielectrics.

We cannot compute $\mathbf{P}$ directly from this equation.

$$\mathbf{E}_0 \rightarrow \mathbf{P}_0$$

$$\mathbf{P}_0 \rightarrow \mathbf{E}_0 + \Delta \mathbf{E}'_P$$

$$\mathbf{E}_0 + \Delta \mathbf{E}'_P \rightarrow \mathbf{P}_0 + \Delta \mathbf{P}'_0$$





What did we learn in Chap. 4?

Now we are going to treat the field caused by both bound charge and free charge.

$$\begin{aligned}\rho &= \rho_f + \rho_b \\ &= \rho_f - \nabla \cdot \mathbf{P} = \epsilon_0 \nabla \cdot \mathbf{E}\end{aligned}$$

where $\mathbf{E}$ is now the **total field**, not just that portion generated by polarization.

$$\epsilon_0 \nabla \cdot \mathbf{E} + \nabla \cdot \mathbf{P} = \rho_f$$

$$\mathbf{D} = \epsilon_0 \mathbf{E} + \mathbf{P} = \epsilon_0 \mathbf{E} + \epsilon_0 \chi_e \mathbf{E} = \epsilon_0 (1 + \chi_e) \mathbf{E} = \epsilon \mathbf{E}$$

$$\nabla \cdot (\epsilon_0 \mathbf{E} + \mathbf{P}) \equiv \nabla \cdot \mathbf{D} = \rho_f$$

$$\epsilon_r = \frac{\epsilon}{\epsilon_0} = 1 + \chi_e \quad \text{Relative permittivity}$$

Electric displacement

$$\nabla \cdot \mathbf{D} = \rho_f \Rightarrow \oint \mathbf{D} \cdot d\mathbf{a} = Q_{f \text{ enc}}$$


$$\mathbf{E} = \frac{1}{\epsilon} \mathbf{D} = \frac{1}{\epsilon_r} \mathbf{E}_{\text{vac}}$$

Space filled with a homogenous linear dielectric $\rightarrow$ field reduced by a factor of one over the dielectric constant
 $\rightarrow$ **Polarization partially shields the charge**



What did we learn in Chap. 4?

Problem 4.18 The space between the plates of a parallel-plate capacitor (Fig. 4.24) is filled with two slabs of linear dielectric material. Each slab has thickness a , so the total distance between the plates is $2a$. Slab 1 has a dielectric constant of 2, and slab 2 has a dielectric constant of 1.5. The free charge density on the top plate is σ and on the bottom plate $-\sigma$.



$$\oint_S \mathbf{D} \cdot d\mathbf{a} = Q_{f_{enc}} \Rightarrow 2DA = \sigma A \Rightarrow \mathbf{D} = \frac{\sigma}{2} \hat{\mathbf{z}}$$

$$\therefore \mathbf{D} = \begin{cases} \mathbf{0}, & \text{outside the plates} \\ -\frac{\sigma}{2} \hat{\mathbf{z}} + \frac{\sigma}{2} \hat{\mathbf{z}} = -\sigma \hat{\mathbf{z}} \end{cases}$$

$$\mathbf{E} = \frac{\mathbf{D}}{\epsilon} \Rightarrow \begin{cases} \mathbf{E}_1 = -\frac{\sigma}{2\epsilon_0} \hat{\mathbf{z}}, & \text{for slab 1} \\ \mathbf{E}_2 = -\frac{2\sigma}{3\epsilon_0} \hat{\mathbf{z}}, & \text{for slab 2} \end{cases} \Rightarrow \Delta V = -\int_-^+ \mathbf{E} \cdot d\vec{\ell} = \frac{7\sigma a}{6\epsilon_0}$$

$$\mathbf{P} = \epsilon_0 \chi \mathbf{E} = \epsilon (\epsilon_0 - 1) \mathbf{E} = \epsilon (\epsilon - 1) \frac{-\sigma}{\epsilon_0} \hat{\mathbf{z}} = -\sigma \left(1 - \frac{1}{\epsilon_r} \right) \hat{\mathbf{z}}$$


$$\therefore \mathbf{P}_1 = -\frac{\sigma}{2} \hat{\mathbf{z}}, \quad \mathbf{P}_2 = -\frac{\sigma}{3} \hat{\mathbf{z}} \Rightarrow \nabla \cdot \mathbf{P} = 0 \Rightarrow \rho_b = 0 \text{ everywhere}$$

	$+\sigma$
①	$-\sigma/2$
	$+\sigma/2$
②	$-\sigma/3$
	$+\sigma/3$
	$-\sigma$






What did we learn in Chap. 4?



(a)



(b)

$$\left(\frac{C_b}{C_0} = \frac{1+\epsilon_r}{2}\right) > \left(\frac{C_a}{C_0} = \frac{2}{1+\epsilon_r^{-1}}\right)$$




	E	D	P
(a) air	$\frac{2\epsilon_r}{(\epsilon_r+1)} \frac{V}{d} \hat{x}$	$\frac{2\epsilon_r}{(\epsilon_r+1)} \frac{\epsilon_0 V}{d} \hat{x}$	0
(a) dielectric	$\frac{2}{(\epsilon_r+1)} \frac{V}{d} \hat{x}$	$\frac{2\epsilon_r}{(\epsilon_r+1)} \frac{\epsilon_0 V}{d} \hat{x}$	$\frac{2(\epsilon_r-1)}{(\epsilon_r+1)} \frac{\epsilon_0 V}{d} \hat{x}$
(b) air	$\frac{V}{d} \hat{x}$	$\frac{\epsilon_0 V}{d} \hat{x}$	0
(b) dielectric	$\frac{V}{d} \hat{x}$	$\epsilon_r \frac{\epsilon_0 V}{d} \hat{x}$	$(\epsilon_r - 1) \frac{\epsilon_0 V}{d} \hat{x}$

	σ_b (top surface)	σ_f (top plate)
(a)	$-\frac{2(\epsilon_r-1)}{(\epsilon_r+1)} \frac{\epsilon_0 V}{d}$	$\frac{2\epsilon_r}{(\epsilon_r+1)} \frac{\epsilon_0 V}{d}$
(b)	$-(\epsilon_r - 1) \frac{\epsilon_0 V}{d}$	$\epsilon_r \frac{\epsilon_0 V}{d}$ (left); $\frac{\epsilon_0 V}{d}$ (right)





What did we learn in Chap. 4?



Ref: *AJP* **73**, 52 (2005)





What did we learn in Chap. 4?

$$\rho_b = -\nabla \cdot \mathbf{P} = -\nabla \cdot \left(\epsilon_0 \chi_e \frac{\mathbf{D}}{\epsilon} \right) = -\frac{\chi_e}{1 + \chi_e} \rho_f \quad \leftarrow \text{in a homogenous linear dielectric}$$

$$\epsilon = \epsilon_0 (1 + \chi_e)$$

shielding effect


$$D_{above}^{\perp} - D_{below}^{\perp} = \sigma_f \quad \Rightarrow \quad \epsilon_{above} E_{above}^{\perp} - \epsilon_{below} E_{below}^{\perp} = \sigma_f$$

$$(\epsilon_{above} \nabla V_{above} - \epsilon_{below} \nabla V_{below}) = -\sigma_f \hat{\mathbf{n}}$$





What did we learn in Chap. 4?



$$\left. \begin{array}{l} \text{(i)} \quad V_{\text{in}} = V_{\text{out}} \quad \text{at } r = R \\ \text{(ii)} \quad \epsilon \frac{\partial V_{\text{in}}}{\partial r} = \epsilon_0 \frac{\partial V_{\text{out}}}{\partial r} \quad \text{at } r = R \\ \text{(iii)} \quad V_{\text{out}} \rightarrow -E_0 r \cos\theta \quad \text{for } r \gg R \end{array} \right\}$$

$$\left\{ \begin{array}{l} V_{\text{in}}(r, \theta) = -\frac{3E_0}{\epsilon_r + 2} r \cos\theta \\ V_{\text{out}}(r, \theta) = -E_0 r \cos\theta + \left(\frac{\epsilon_r - 1}{\epsilon_r + 2} \right) R^3 E_0 r^{-2} \cos\theta \end{array} \right.$$


$$\mathbf{E}_{\text{in}} = -\nabla V_{\text{in}} = \frac{3E_0}{\epsilon_r + 2} \hat{\mathbf{z}} \quad \leftarrow \text{uniform}$$

$$V(r, \theta) = \sum_{\ell=0}^{\infty} (A_{\ell} r^{\ell} + B_{\ell} r^{-(\ell+1)}) P_{\ell}(\cos\theta)$$

$$\left\{ \begin{array}{l} V_{\text{in}}(r, \theta) = \sum_{\ell=0}^{\infty} A_{\ell} r^{\ell} P_{\ell}(\cos\theta) \quad r \leq R \\ V_{\text{out}}(r, \theta) = -E_0 r \cos\theta + \sum_{\ell=0}^{\infty} B_{\ell} r^{-(\ell+1)} P_{\ell}(\cos\theta) \quad r \geq R \end{array} \right.$$


How to connect this result with Polarization?

$$\mathbf{P} = \epsilon_0 \chi_e \mathbf{E}$$





What did we learn in Chap. 4?



$$\mathbf{E}_0 \rightarrow \mathbf{P}_0 = \epsilon_0 \chi_e \mathbf{E}_0 \rightarrow \mathbf{E}_1 = \frac{-\mathbf{P}_0}{3\epsilon_0} = \frac{-\chi_e}{3} \mathbf{E}_0$$

$$\mathbf{E}_1 \rightarrow \mathbf{P}_1 = \epsilon_0 \chi_e \mathbf{E}_1 = \frac{-\epsilon_0 \chi_e^2}{3} \mathbf{E}_0 \rightarrow \mathbf{E}_2 = \frac{-\mathbf{P}_1}{3\epsilon_0} = \frac{\chi_e^2}{9} \mathbf{E}_0$$

$\vdots$


$$\mathbf{E}_n = \left(\frac{-\chi_e}{3} \right)^n \mathbf{E}_0$$

$$\therefore \mathbf{E} = \mathbf{E}_0 + \mathbf{E}_1 + \mathbf{E}_2 + \dots = \sum_{n=0}^{\infty} \left(\frac{-\chi_e}{3} \right)^n \mathbf{E}_0 = \frac{\mathbf{E}_0}{1 + (\chi_e / 3)} = \frac{3\mathbf{E}_0}{\epsilon_r + 2}$$

Remark: 1. Consistent with the previous result
2. This method formally requires $\chi_e < 3$



What did we learn in Chap. 4?




Partial Image Charge

$$W = \frac{1}{2} \int (\epsilon_0 \mathbf{E} \cdot \mathbf{E}) d\tau$$

$$W = \frac{1}{2} \int (\mathbf{E} \cdot \mathbf{D}) d\tau$$

Energy in Dielectric systems



Force on Dielectric
Fringing Field Effect



What did we learn in Chap. 4?

In a linear dielectric, the polarization is said to be proportional to the field $\mathbf{P} = \epsilon_0 \chi_e \mathbf{E}_{macro}$ $\leftarrow$ Macroscopic $\mathbf{E}_{macro} = \mathbf{E}_{self} + \mathbf{E}_{else}$

If the material consists of atoms (or nonpolar molecules), the induced dipole moment $\mathbf{p} = \alpha \mathbf{E}_{else}$ $\leftarrow$ Microscopic

$$\mathbf{P} = N\mathbf{p} = N\alpha \mathbf{E} \stackrel{?}{\Rightarrow} \chi_e = \frac{N\alpha}{\epsilon_0}$$

If the density of atoms is low, it's not far off. However, the fields used are from different viewpoints!

$$\mathbf{E}_{self} = \frac{-\mathbf{P}}{4\pi\epsilon_0 R^3} \Rightarrow \mathbf{E}_{macro} = \frac{-\alpha}{4\pi\epsilon_0 R^3} \mathbf{E}_{else} + \mathbf{E}_{else} = \left(1 - \frac{N\alpha}{3\epsilon_0}\right) \mathbf{E}_{else} = \frac{\mathbf{P}}{\epsilon_0 \chi_e} = \frac{N\alpha}{\epsilon_0 \chi_e} \mathbf{E}_{else}$$

$$\therefore \chi_e = \frac{N\alpha\epsilon_0}{1 - N\alpha\epsilon_0} \Rightarrow \alpha = \frac{3\epsilon_0}{N} \frac{\chi_e}{3 + \chi_e} = \frac{3\epsilon_0}{N} \frac{\epsilon_r - 1}{\epsilon_r + 2} \approx \frac{3\epsilon_0}{N} \frac{n^2 - 1}{n^2 + 2}$$

What about polar substance?

Clausius-Mossotti formula

Lorentz-Lorenz relation





What did we learn in Chap. 4?

Energy of a dipole in an external field

$$u = -\mathbf{p} \cdot \mathbf{E} = -pE \cos\theta$$

Statistical mechanics says that for a material in equilibrium at absolute temperature, the probability of a given molecule having energy is proportional to the Boltzmann factor

$$\exp(-u/kT)$$

The average energy of the dipoles is therefore

$$\langle u \rangle = \frac{\int u e^{-u/kT} d\Omega}{\int e^{-u/kT} d\Omega} = \frac{\int_{-pE}^{pE} u e^{-u/kT} du}{\int_{-pE}^{pE} e^{-u/kT} du} = kT - pE \left[\frac{e^{pE/kT} + e^{-pE/kT}}{e^{pE/kT} - e^{-pE/kT}} \right] = kT - pE \coth(pE/kT)$$


Linear region

$$P \approx \frac{Np^2}{3kT} E = \epsilon_0 \chi_e E$$

$$\Rightarrow \chi_e = \frac{Np^2}{3\epsilon_0 kT}$$

$$\therefore \|\mathbf{P}\| = N \|\langle \mathbf{p} \rangle\| = N \left\| \langle \mathbf{p} \cdot \mathbf{E} \rangle \frac{\hat{\mathbf{E}}}{E} \right\| = -Np \frac{\langle u \rangle}{pE} = Np \left\{ \coth\left(\frac{pE}{kT}\right) - \frac{kT}{pE} \right\}$$

Langevin equation



Comment: For large fields/low temperatures, all the molecules are lined up, and the material is nonlinear.



Remarks on Dielectrics

Anisotropic Materials:


i-th component of the polarization is related to the j-th component of the electric field.

$$P_i = \sum_j \epsilon_0 \chi_{ij} E_j$$

Polarize in the x direction by applying a field in the z direction → Crystal Optics.

How to choose “unit volume” in reality?
e.g., Plasma in **microscopic** scale is regarded as **a gas of free charges** ($P = 0$). However, in **macroscopic** scale, it serves as **a continuous medium**, exhibiting **non-zero permittivity**.

Polarization Ambiguity



Non-uniqueness of P is not problematic, because **every measurable consequence of P is in fact a consequence of a continuous change in P .**

